



## Shoulder Arthroplasty Postoperative Instructions

### General

A family member or caregiver may need to assist you for the next few weeks.

- Helps with basic movement and function
- Helps with household chores
- Helps ensure all medications are administered
- Helps provide emotional support for daily life and rehabilitation

Furniture may need to be rearranged to avoid tripping and falling.

In general, the more you are up and ambulating, the better you will do.

You may ice once every hour for up to 20 minutes, as needed, in the first few postoperative days.

If you had a nerve block, the effects may last 24 hours.

### Activity

#### **Do:**

- Wear the sling at all times, except for showering or while getting dressed. This includes wearing the sling while sleeping.
- You may come out of the sling to perform therapy exercises at home. Your therapist will show you these exercises.
- Perform gentle elbow, wrist, finger motion exercises several times per day.
- Follow your physical therapist's instructions.
- You may return to work with minimal use of the operative arm (ie desk work)

#### **Do Not:**

- Lift anything with the operative arm

### Wound Care

Unless otherwise noted, your surgical site was closed with all-absorbable sutures under the skin, with glue over top to seal the incision. A clear dressing was placed over top in the operating room.

- Leave the dressing in place. Dr. Baessler will remove it at your postoperative follow up appointment.

A small amount of drainage is normal for the first few days postoperatively.

- Excessive drainage that leaks out from under the dressing is worrisome. Please call the office with any questions or concerns.

Some redness around the incision is also normal in the first week postoperatively.



- If the wound becomes progressively more red, swollen, or painful after several days, please call the office.

## Medications

Unless otherwise noted, you will be given prescriptions for the following medications:

Aspirin 81 mg daily (2 weeks total)

- Helps prevent blood clots postoperatively

Acetaminophen (Tylenol) 1000 mg every 8 hours (4 weeks total)

- This high dose, combined with gabapentin and celecoxib, works very well for pain without the need for opioids.
- Do not exceed 3000 mg in a single day

Gabapentin (Neurontin) 100 mg three times daily (4 weeks total)

- Works along with acetaminophen and celecoxib
- May make you drowsy

Celecoxib (Celebrex) 100 mg twice daily (4 weeks total)

- Works along with acetaminophen and gabapentin
- Stay hydrated while on this medication

Oxycodone (Roxicodone) 5 mg every 4 hours as needed (1 week total)

- Strong pain medication that can cause addiction
- Indiana law only allows prescription of 1 weeks' worth of narcotics
- Only use this as a last resort
- May cause nausea, constipation, drowsiness, altered mentation

Docusate (Colace) 100 mg twice daily (2 weeks total)

- This helps prevent constipation associated with oxycodone

Ondansetron (Zofran) 4 mg dissolved on tongue every 6 hours as needed for nausea

- For postoperative nausea

## Showering

You may shower on postoperative day 3

- Leave the dressing in place. It seals the incision.
- Let water run over the incision. Do not vigorously scrub. Pat dry.

Do NOT soak or submerge under water (bath tub, hot tub, pool).

Aaron M. Baessler, MD  
Orthopaedic Surgery  
Sports Medicine, Shoulder and Elbow Surgery

Muncie Office  
3631 N. Morrison Rd  
Muncie, IN 47304  
(317) 208-3866



Franklin Office  
990 East State Rd 44  
Franklin, IN 46131  
(317) 208-3866

- Until wound is completely healed (usually around 6 weeks)

### Driving

You may return to driving if:

- You are completely off narcotic pain medications
- Strength and reflexes are back to normal
- Typically no earlier than 2 weeks postoperatively

### Follow-up

Dr. Baessler will see you back in the office 2 weeks postoperatively for a wound check.